

The Nuclear Level Density and the Determination of Thermonuclear Rates for Astrophysics

T. RAUSCHER¹, F.-K. THIELEMANN¹, K.-L. KRATZ²

¹*Institut für theoretische Physik, Universität Basel, Basel, Switzerland*

²*Institut f. Kernchemie, Universität Mainz, Mainz, Germany*

ABSTRACT. The prediction of cross sections for nuclei far off stability is crucial in the field of nuclear astrophysics. In recent calculations the nuclear level density – as an important ingredient to the statistical model (Hauser-Feshbach) – has shown the highest uncertainties. We present a global parametrization of nuclear level densities based on the back-shifted Fermi-Gas formalism. Employment of an energy-dependent level density parameter a and microscopic corrections from a recent FRDM mass formula by Möller et al. leads to a highly improved fit of level densities at the neutron-separation energy in the mass range $20 \leq A \leq 245$. The importance of using proper microscopic corrections from mass formulae is emphasized. The resulting level description is well suited for astrophysical applications.

The level density can also provide clues to the applicability of the statistical model which is only correct for a high density of excited states. Using the above description one can derive a “map” for the applicability of the model for reactions of stable and unstable nuclei with neutral and charged particles.

1. Introduction

Explosive nuclear burning in astrophysical environments produces unstable nuclei, which again can be targets for subsequent reactions. In addition, it involves a very large number of stable nuclei, which are not fully explored by experiments. Thus, it is necessary to be able to predict reaction cross sections and thermonuclear rates with the aid of theoretical models. Explosive burning in supernovae involves in general intermediate mass and heavy nuclei. Due to a large nucleon number they have intrinsically a high density of excited states. A high level density in the compound nucleus at the appropriate excitation energy allows to make use of the statistical model approach for compound nuclear reactions (e.g. Hauser & Feshbach 1952, Mahaux & Weidenmüller 1979), which averages over resonances.

As the capture of an alpha particle leads usually to larger Q -values than neutron or proton captures, the compound nucleus is created at a higher excitation energy and especially in the case of alpha-captures it is often even possible to apply the Hauser-Feshbach formalism for light nuclei. Another advantage of alpha-captures is that the capture Q -values vary little with the N/Z -ratio of a nucleus, for nuclei with $Z \leq 50$. For $Z > 50$, entering the regime of natural alpha-decay, very small alpha-capture Q -values can be encountered for proton-rich nuclei. Such nuclei on the other hand do not play a significant role in astrophysical environments, maybe with exception of the p -process. This means that in the case of alpha-captures the requirement of large level densities

at the bombarding energy is equally well fulfilled at stability as for unstable nuclei. Opposite to the behavior for alpha-induced reactions, the reaction Q-values for proton or neutron captures vary strongly with the N/Z-ratio, leading eventually to vanishing Q-values at the proton or neutron drip line. For small Q-values the compound nucleus is created at low excitation energies and also for intermediate nuclei the level density can be quite small. Therefore, it is not advisable to apply the statistical model approach close to the proton or neutron drip lines for intermediate nuclei. For neutron captures close to the neutron drip line in r-process applications it might be still permissible for heavy and often deformed nuclei, which have a high level density already at very low excitation energies.

In astrophysical applications usually different points are emphasized than for investigations in pure nuclear physics. Many of the latter in this well established field were focused on specific reactions, where all or most "ingredients", like optical potentials for particle and alpha transmission coefficients, level densities, resonance energies and widths of giant resonances to be implemented in predicting E1 and M1 gamma-transitions, were deduced from experiments. This of course, as long as the statistical model prerequisites are met, will produce highly accurate cross sections. For the majority of nuclei in astrophysical applications such information is not available. The real challenge is thus not the well established statistical model, but rather to provide all these necessary ingredients in as reliable a way as possible, also for nuclei where none of such informations are available. In addition, these approaches should be on a similar level as e.g. mass models, where the investigation of hundreds or thousands of nuclei is possible with manageable computational effort, which is not always the case for fully microscopic calculations.

The major uncertainty in all existing calculations stems from the prediction of nuclear level densities (Truran et al. 1966; Holmes et al. 1976; Cowan, Thielemann & Truran 1991, and extended references therein), which in earlier calculations gave uncertainties even beyond a factor of 10 at the neutron separation energy (Gilbert & Cameron 1965), about a factor of 8 (Woosley et al. 1978), and a factor of 5 even in the most recent calculations (e.g. Thielemann, Arnould & Truran 1987; see also Fig. 3.16 in Cowan, Thielemann & Truran 1991).

2. Thermonuclear Rates from Statistical Model Calculations

A high level density in the compound nucleus permits to use averaged transmission coefficients T , which do not reflect a resonance behavior, but rather describe absorption via an imaginary part in the (optical) nucleon-nucleus potential (for details see Mahaux & Weidenmüller 1979). This leads to the well known expression

$$\sigma_i^{\mu\nu}(j, o; E_{ij}) = \frac{\pi\hbar^2/(2\mu_{ij}E_{ij})}{(2J_i^\mu + 1)(2J_j + 1)} \times \sum_{J, \pi} (2J + 1) \frac{T_j^\mu(E, J, \pi, E_i^\mu, J_i^\mu, \pi_i^\mu) T_o^\nu(E, J, \pi, E_m^\nu, J_m^\nu, \pi_m^\nu)}{T_{tot}(E, J, \pi)} \quad (1)$$

for the reaction $i^\mu(j, o)m^\nu$ from the target state i^μ to the excited state m^ν of the final nucleus, with center of mass energy E_{ij} and reduced mass μ_{ij} . J denotes the spin, E the excitation energy, and π the parity of excited states. When these properties are used without subscripts they describe the compound nucleus, subscripts refer to states of the participating nuclei in the reaction $i^\mu(j, o)m^\nu$ and superscripts indicate the specific excited states. Experiments measure $\sum_\nu \sigma_i^{0\nu}(j, o; E_{ij})$, summed over all excited states of the final nucleus, with the target in the ground state. Target states μ in an astrophysical plasma are thermally populated and the astrophysical cross section $\sigma_i^*(j, o)$ is given by

$$\sigma_i^*(j, o; E_{ij}) = \frac{\sum_\mu (2J_i^\mu + 1) \exp(-E_i^\mu/kT) \sum_\nu \sigma_i^{\mu\nu}(j, o; E_{ij})}{\sum_\mu (2J_i^\mu + 1) \exp(-E_i^\mu/kT)} \quad (2)$$

The summation over ν replaces $T_o^\nu(E, J, \pi)$ in Eq. 1 by the total transmission coefficient

$$T_o(E, J, \pi) = \sum_{\nu=0}^{\nu_m} T_o^\nu(E, J, \pi, E_m^\nu, J_m^\nu, \pi_m^\nu) + \int_{E_m^{\nu_m}}^{E-S_{m,o}} \sum_{J_m, \pi_m} T_o(E, J, \pi, E_m, J_m, \pi_m) \rho(E_m, J_m, \pi_m) dE_m \quad (3)$$

Here $S_{m,o}$ is the channel separation energy, and the summation over excited states above the highest experimentally known state ν_m is changed to an integration over the level density ρ . The summation over target states μ in Eq. 2 has to be generalized accordingly.

In addition to the ingredients required for Eq. 1, like the transmission coefficients for particles and photons, the width fluctuation corrections $W(j, o, J, \pi)$ have to be employed. The important ingredients of statistical model calculations as indicated in Eqs. 1 through 3 are the particle and γ -transmission coefficients T and the level density of excited states ρ . Therefore, the reliability of such calculations is determined by the accuracy with which these components can be evaluated.

3. Level Densities

3.1. The Back-Shifted Fermi-Gas Model

Considerable effort has been put into the improvement of the input for the statistical Hauser-Feshbach calculations (e.g. Cowan, Thielemann & Truran 1991). However, the nuclear level density has given rise to the largest uncertainties in the description of nuclear reactions. For calculating the level densities in that context one does not only have to find reliable methods, but also computationally feasible ones. In dealing with thousands of nuclei one has to resort to simple models in order to minimize computer time.

Such a simple model is the non-interacting Fermi-gas model introduced by Bethe (1936). Mostly, the back-shifted Fermi-gas description, assuming an even distribution of odd and even parities, is used (Gilbert & Cameron 1965):

$$\rho(U, J, \pi) = \frac{1}{2} f(U, J) \rho(U) \quad (4)$$

with

$$\begin{aligned} \rho(U) &= \frac{1}{\sqrt{2\pi}} \frac{\sqrt{\pi}}{12a^{1/4}} \frac{\exp(2\sqrt{aU})}{U^{5/4}} , & f(U, J) &= \frac{2J+1}{2\sigma^2} \exp\left(\frac{-J(J+1)}{2\sigma^2}\right) \\ \sigma^2 &= \frac{\Theta_{\text{rigid}}}{\hbar^2} \sqrt{\frac{U}{a}} , & \Theta_{\text{rigid}} &= \frac{2}{5} m_u A R^2 , & U &= E - \delta \quad . \end{aligned} \quad (5)$$

The spin dependence is determined by the spin cut-off parameter σ . Thus, the level density is dependent on only two parameters: the level density parameter a and the backshift δ , which determines the energy of the first excited state.

Within this framework, the quality of level density predictions depends on the reliability of systematic estimates of a and δ . Gilbert & Cameron (1965) were the first to identify an empirical correlation of the level density parameter a with experimental shell corrections $S(N, Z)$ and of the backshift δ with experimental pairing corrections.

An improved approach has to consider the energy dependence of the shell effects which are known to vanish at high excitation energies (Iljinov et al. 1992), i.e. the thermal damping of shell effects. Although, for astrophysical purposes only energies close to the particle separation thresholds have to be considered, an energy dependence can lead to a considerable improvement of the global fit. This is especially true for strongly bound nuclei close to magic numbers.

We use the following description (initially proposed by Ignatyuk, Smirenkin & Tishin 1975; Ignatyuk, Istekov & Smirenkin 1979) for an energy-dependent level density parameter a :

$$a(U, Z, N) = \tilde{a}(A) \left[1 + C(Z, N) \frac{f(U - \delta)}{U - \delta} \right] , \quad (6)$$

where

$$\tilde{a}(A) = \alpha A + \beta A^{2/3} \quad (7)$$

and

$$f(U) = 1 - \exp(-\gamma U) \quad . \quad (8)$$

The shape of the function $f(U)$ was found by approximation of numerical microscopic calculations based on the shell model. Thus, we are left with three open parameters, namely α , β , and γ . The values of these parameters can be determined by fitting experimental data.

Important input for these fits are the so-called shell correction $C(Z, N)$ and the pairing gap Δ . These have to be taken from microscopic or macroscopic-microscopic nuclear mass formulae. Instead of assuming constant pairing or a fixed dependence on the mass number A , we directly determine the pairing gap Δ from mass differences of neighboring nuclei. Thus, for the neutron pairing gap Δ_n one obtains (Wang et al. 1992)

$$\Delta_n(Z, N) = \frac{1}{4} [E^G(Z, N - 2) - 3E^G(Z, N - 1) + 3E^G(Z, N) - E^G(Z, N + 1)] \quad , \quad (9)$$

where $E^G(Z, N)$ is the mass for the nucleus (Z, N) . Similarly, the proton pairing gap Δ_p can be calculated.

Previous attempts to find a global description of level densities used shell corrections derived from comparison of liquid-drop masses with experiment ($C = S \equiv M_{\text{exp}} - M_{\text{LD}}$) or the “empirical” shell corrections $C(Z, N) = S(Z, N)$ given by Gilbert & Cameron (1965). A problem connected with the use of liquid-drop masses arises from the fact that there are different liquid-drop model parametrizations available in the literature which produce quite different values for S .

We realized that one has to reconsider the meaning of the parameter C in the level density formula (Eq. 6). Actually, it should describe properties of a nucleus differing from the *spherical* macroscopic energy (i.e. mass), including properties which are vanishing at higher excitation energies. Therefore, the parameter C should rather be identified with the so-called “microscopic” correction E_{mic} . The mass of a nucleus with deformation ϵ can then be written as

$$M_{\text{theo}}(\epsilon) = E_{\text{mic}}(\epsilon) + E_{\text{mac}}(\text{sphere}) \quad . \quad (10)$$

Alternatively, one can write

$$M_{\text{theo}}(\epsilon) = E_{\text{mac}}(\epsilon) + E_{\text{s+p}}(\epsilon) \quad , \quad (11)$$

with $E_{\text{s+p}}$ being the shell-plus-pairing correction. The confusion about the term “microscopic correction”, which is sometimes ambiguously used in literature, is also pointed out in Möller et al. (1995). Thus, the above mentioned ambiguity of different liquid-drop parametrizations follows from the inclusion of partially deformation dependent effects into the macroscopic part of the mass formula.

3.2. Results

In our study we utilized the most recent mass formula by Möller et al. (1995), consistently also taking the microscopic corrections of that mass formula (calculated in a folded Yukawa shell model with Lipkin-Nogami pairing) in order to determine the parameter $C(Z, N) = E_{\text{mic}}$. The backshift δ was calculated by setting $\delta(Z, N) = \Delta_{\text{n}}(Z, N) + \Delta_{\text{p}}(Z, N)$ and using Eq. 9. In order to obtain the parameters α , β , and γ , we performed a fit to experimental data on s-wave neutron resonance spacing of 278 nuclei at the neutron separation energy. The data were taken from the compilation by Iljinov et al. (1992).

As a quantitative overall estimate of the agreement between calculation and experiment, one usually quotes the averaged ratio

$$g \equiv \left\langle \frac{\rho_{\text{calc}}}{\rho_{\text{exp}}} \right\rangle = \exp \left[\frac{1}{n} \sum_{i=1}^n \left(\ln \frac{\rho_{\text{calc}}^i}{\rho_{\text{exp}}^i} \right)^2 \right]^{1/2} \quad , \quad (12)$$

with n being the number of nuclei for which level densities ρ are experimentally known.

As a best fit (when including known experimental masses where known), we obtain an averaged ratio $g = 1.5$ with the parameter values $\alpha = 0.1336$, $\beta = -0.06712$, $\gamma = 0.04862$. The ratios of experimental to predicted level densities for the nuclei considered is shown in Fig. 1. As can be seen, for the majority of nuclei the absolute deviation is less than a factor of 2. This is a quite satisfactory improvement over the theoretical level

densities used previously. Such a direct comparison was rarely shown in earlier work, mostly the level density parameter a , entering exponentially into the level density, was displayed. Closely examining the nuclei with the largest deviations in our fit, we were not able to find any remaining correlation of the deviation with separation energy (i.e. excitation energy) or spin. Different combinations of masses and microscopic corrections from other models (droplet model by Myers and Swiatecki, Cameron-Elkin mass formula and shell corrections, experimental values) were also tried but did not lead to better results.

As the parameters were fit to experimentally accessible nuclei close to the drip line, the question arises how reliable such a level density description can be for astrophysically interesting nuclei far from stability. Since the microscopic corrections (and the masses) are taken from a nuclear-structure model this leads to the question of the reliability of that model far off stability. There is, of course, no final conclusion regarding that point yet. However, recent investigations in astrophysics and nuclear physics have shown the robustness of the Möller et al. (1995) approach (Thielemann et al. 1994). Recently improved purely microscopic models have exhibited similar behavior towards the drip lines (Dobaczewski et al. 1994), but there are no large scale calculations over the whole chart of nuclei available yet. Therefore, the model used here is among the most reliable ones available at present.

4. Applicability of the Statistical Model

With the help of a level density description one is also able to make statements on when and where the statistical model approach is valid. Generally speaking, in order to apply the model correctly a sufficiently large number of levels is needed in the relevant energy range which can act as doorway states to form a compound nucleus. In the following we will discuss this with the aid of the above level density approach for neutron-, proton- and α -induced reactions. This section is intended to be a guide to a meaningful and correct application of the statistical model.

4.1. Neutron-Induced Reactions

In the case of neutron-induced reactions a criterion for the applicability can directly be derived from the level density. For astrophysical purposes the projectile energies are quite low by nuclear physics standards. Therefore, the relevant energies will lie very close to the neutron separation energy. Thus, one just has to consider the level density at this energy. As a rule-of-thumb it is usually said that there should at least be 10 levels per MeV for reliable statistical model calculations. The level densities at the appropriate neutron separation energies are shown in Fig. 2 (note that therefore the level density is plotted at a *different* energy for each nucleus). One can easily identify the magic neutron numbers by the drop in level density, as well as odd-even staggering effects. A general sharp drop is also found for nuclei close to the neutron drip line. For nuclei with such low level densities the statistical model cross sections will become very small and other processes might become important, such as direct reactions.




Fig. 1. Ratio of predicted to experimental (Iljinov et al. 1992) level densities at the neutron separation energy.

Level Density at S_n




Fig. 2. Level density (levels per MeV) at the neutron separation energy.

4.2. Charged-Particle Induced Reactions

Contrary to neutron-induced reactions, the knowledge of the level density at the particle separation energy is not sufficient to say something about the reliability of the statistical model because in this case also the Coulomb barrier has to be taken into account. By folding the Maxwell-Boltzmann velocity distribution of the projectiles at a given temperature with the penetrability through the Coulomb barrier, the so-called Gamow peak can be derived, in which most of the reactions will take place. Location and width of the Gamow peak depend on the charges of projectile and target, and on the temperature.

Similar to the rule-of-thumb given above for neutron-induced reactions, an estimate can be made of how many resonances have to be inside the Gamow peak in order to still be able to numerically solve the integral when calculating the reaction rate. Comparison with experimental cross sections gives the result that one can assume a value of about 3 resonances (levels). However, there are comparisons to experimental data (Van Wormer et al. 1994) which have shown that in certain cases the experimental values could be reproduced with even fewer levels. Nevertheless, to be on the safe side we assumed 3.5 levels within the width (FWHM) of the Gamow peak as the necessary minimum in the plots shown in Figs. 3 and 4.

For proton-induced reactions Fig. 3 should be used, whereas Fig. 4 is for α -induced reactions. These plots do not directly display level densities but rather show (again in a logarithmic scale) the temperatures at which the statistical model can be used. That means that for each nucleus (it is always the compound nucleus which is plotted for the reaction) the temperature is given beyond which the above described condition of 3.5 levels is fulfilled. This enables one to directly read from the plot whether the statistical model cross section can be “trusted” or whether single resonances or other processes (e.g. direct reactions) have also to be considered. (However, this does not necessarily mean that the statistical cross section is always negligible in the latter cases, since the assumed condition is quite conservative).

5. Summary

We were able to improve considerably the prediction of nuclear level densities by employing an energy-dependent description for the level density parameter a and by correctly including microscopic corrections. All nuclei can now be described with a single parameter set consisting of just three parameters. The globally averaged deviation of prediction from experiment of about 1.5 translates into somewhat smaller deviations of the final cross sections due to the dominance of transitions to states with low excitation energies. This will also make it worthwhile to recalculate the cross sections and thermonuclear rates for many astrophysically important reactions in the intermediate and heavy mass region.

We also presented a “map” as a guide for the application of the statistical model for neutron-, proton- and α -induced reactions. The above plots can give hints on when it is safe to use the statistical model approach and which nuclei have to be treated with special attention at a given temperature. Thus, information on which nuclei might be of special interest for an experimental investigation may also be extracted. It should be

Applicability of the Statistical Model (p)




Fig. 3. Temperatures (in T_g) for which the statistical model can be used. Plotted is the compound nucleus of the proton-induced reaction $p + \text{Target}$ (see text).




Fig. 4. Temperatures (in T_g) for which the statistical model can be used. Plotted is the compound nucleus of the α -induced reaction $\alpha + \text{Target}$ (see text).

noted that we used very conservative assumptions in deriving the above criteria for the applicability of the statistical model.

Acknowledgements

This work was supported in part by the Swiss Nationalfonds. TR is acknowledging support by an APART fellowship from the Austrian Academy of Sciences.

References

- Bethe, A.H.: 1936, *Phys. Rev.* **50**, 332.
- Dobaczewski, J., Hamamoto, I., Nazarewicz, W., Sheikh, J.A.: 1994, *Phys. Rev. Lett.* **72**, 981.
- Gilbert, A., Cameron, A.G.W.: 1965, *Can. J. Phys.* **43**, 1446.
- Hauser, W., Feshbach, H.: 1952, *Phys. Rev.* **A87**, 366.
- Holmes, J.A., Woosley, S.E., Fowler, W.A., Zimmerman, B.A.: 1976, *At. Data Nucl. Data Tables* **18**, 306.
- Ignatyuk, A.V., Smirenkin, G.N., Tishin, A.S.: 1975, *Yad. Phys.* **21**, 485.
- Ignatyuk, A.V., Istekov, K.K., Smirenkin, G.N.: 1979, *Sov. J. Nucl. Phys.* **29**, 450.
- Iljinov, A.S., et al.: 1992, *Nucl. Phys.* **A543**, 517.
- Mahaux, C., Weidenmüller, H.A.: 1979, *Ann. Rev. Part. Nucl. Sci.* **29**, 1.
- Möller, P., Nix, J.R., Myers, W.D., and Swiatecki, W.J.: 1995, *At. Data Nucl. Data Tables* **59**, 185.
- Thielemann, F.-K., Arnould, M., Truran, J.W.: 1987, in *Advances in Nuclear Astrophysics*, eds. E. Vangioni-Flam et al., (Editions frontières: Gif sur Yvette), p. 525.
- Thielemann, F.-K., Kratz, K.-L., Pfeiffer, B., Rauscher, T., van Wormer, L., Wiescher, M.: 1994, *Nucl. Phys.* **A570**, 329c.
- Truran, J.W., Cameron, A.G.W., Gilbert, A.: 1966, *Can. J. Phys.* **44**, 563.
- Van Wormer, L., Görres, J., Iliadis, C., Wiescher, M., Thielemann, F.-K.: 1994, *Astrophys. J.* **432**, 326.
- Wang, R.-P., Thielemann, F.-K., Feng, D.H., Wu, C.-L.: 1992, *Phys. Lett.* **B284**, 196.
- Woosley, S.E., Fowler, W.A., Holmes, J.A., Zimmerman, B.A.: 1978, *At. Nucl. Data Tables* **22**, 371.